



PURPOSE

- Evidence-Based Practice Attitude Scale (EBPAS) measures professionals' feelings about using evidence-based interventions
- Use of evidence-based interventions is novel in Brazilian mental health settings

Goal: translate and validate EBPAS for use in Brazilian mental health settings

METHODS

Procedure


- Translated the EBPAS-15 into Brazilian Portuguese; evaluated forward and backtranslations
- Participants recruited via snowball sample
- Collected online August 2019 January 2020
- Performed confirmatory factor analysis (CFA) to determine if scale retained original structure
- Missing values imputed with Random Forest algorithm
- Conducted qualitative interviews examining cultural relevance of items

Sample

- N = 352, mostly:
- Women
- Studied psychology
- Worked as therapist
- •8-14 years of experience
- Worked in private practice

RESULTS

The original factor structure required minor modification.



Ethics laws in Br allow for particip incentives

Most psychologis practice, present for reach


Translation and Validation of the Evidence-Based Practice Attitude Scale to Brazilian Portuguese: Challenges and Lessons Learned

¹Washington University in St. Louis ²Utah State University ³Secretaria de Estado de Desenvolvimento Social Sedes – GDF ⁴Lesley University

allenges	Solutions	
Brazil do not ipant monetary	Include different recruitment strategies such as WhatsApp	
gists are in private nting a challenge	Connect with national organizations (e.g., National Psychology Association) for sampling	

⁵Eastern Illinois University

- ⁶Centro Universitário UniCEUB
- ⁷Centro Paradigma de Análise do Comportamento
- ⁸Aiutare Instituto de Psicologia



"De maneira geral as considero inflexíveis com pouco espaço para acolher e ouvir os pais, o que acaba sendo uma demanda importante deles nesse tipo de rabalho." [Generally, I consider them (the nterventions) not so flexible and with little space to welcome and listen to parents, which ends up being an important demand n this type of work.]

1 Factor 4 Facto 4 Factor * p < 0.05

Washington University in St. Louis

Ana A. Baumann¹ Alejandro J.Vázquez² Bobbi J. Carothers¹ Acileide C.F. Coelho³ Alice Lima Custodio⁴ Mariana M. Juras⁵ Mariana Kohlsdorf⁶ Ana Carolina Macchione⁷ Michela Rodrigues Ribeiro⁸

Summary Statistics for Complete Cases for EBPAS Scale and Items (n = 282)

Items	M, SD	Alpha	
red	2.97, 0.89	0.87	
was required by your supervisor/administrator?	3.10, 0.96		
was required by your school?	3.00, 0.99		
was required by your state?	2.80, 1.04		
	3.43, 0.69	0.73	
was intuitively appealing?	3.48, 1.05		
"it made sense" to you?	3.96, 0.90		
was being used by colleagues who were happy with it?	3.23, 0.91		
ou felt you had enough training to use it correctly?	3.05, 0.81		
ness	3.48, 0.74	0.77	
ke to use new types of methods/interventions to help clients	3.95, 0.91		
Im willing to try new types of methods/interventions even if I	3.69, 1.02		
to follow a teaching/training manual			
im willing to use new types of methods/interventions	3.04, 0.80		
loped by researchers.			
vould try new methods/interventions even if it were very	3.23, 0.99		
rent from what I am used to doing			
gence	1.99, 0.63	0.60	
now better than academic researchers how to care for clients	2.27, 0.97		
esearch-based teaching methods/Interventions are not useful	1.49, 0.94		
actice			
ofessional experience is more important than using	2.44, 1.04		
ualized methods/interventions			
vould not use manualized methods/interventions.	1.77, 0.77	/	

	χ^2	df	χ²/df	RMSEA [90% CI]	SRMR	CFI	TLI
or 15 Items	1054.23	90	11.71	0.175 [0.164 – 0.184] *	0.136	0.904	0.888
ors 15 Items	256.57	84	3.05	0.077 [0.066 – 0.087] *	0.069	0.983	0.978
ors 14 Items	217.17	71	3.06	0.077 [0.065 – 0.088] *	0.066	0.985	0.981

Reference

Aarons, G. A. (2004). Mental health provider attitudes toward adoption of evidence-based practice: The Evidence-Based Practice Attitude Scale (EBPAS). Mental health services research, 6(2), 61-74.

For more information, contact: Ana A. Baumann abaumann@gwbmail.wustl.edu Acknowledgment: This project was funded by the Center for Dissemination and Implementation from the Institute of Public Health, Washington University in St. Louis